



# Ruminal fermentation characteristics and related feeding values of compound feeds and their constituting single feeds studied by using *in vitro* techniques

G. Grubješić , N. Titze<sup>†</sup> , J. Krieg and M. Rodehutschord

Institut für Nutztierwissenschaften, Universität Hohenheim, Emil-Wolff-Str. 6-10, 70599 Stuttgart, Germany

(Received 18 September 2019; Accepted 7 April 2020; First published online 8 May 2020)

*Single concentrate feeds are mixed together forming compound feeds for cattle. However, knowledge regarding the potential interactions (associative effects) between the feeding values of single feeds in compound feeds is lacking. The main objective of the present study was to evaluate ruminal fermentation characteristics and feeding values of eight industrially produced compound feeds in mash form from their constituent single feeds for dairy cows through in vitro assays. Additivity was given for gas production (GP), digestibility of organic matter (dOM) and utilisable CP at the duodenum (uCP). Additivity of CP fractions (determined using the Cornell Net Carbohydrate and Protein System (CNCPS)) was dependent on the fraction and compound feed type; however, the effective degradation calculated from CP fractions (ED<sub>CNCPS</sub>) showed additivity. Additivity was not given for intestinal digestibility of rumen-undegraded protein (ID<sub>RUP</sub>) for five out of eight compound feeds. Precise calculation of metabolisable energy (ME) of compound feeds from ME of single feeds was possible when using the same ME equations for all single and compound feeds. Compound feeds are often provided in pellet form; therefore, our second objective was to evaluate the effects of pelleting on ruminal fermentation characteristics and feeding values of compound feeds. Pelleting affected GP at 24 h (GP<sub>24</sub>; up to 2.4 ml/200 mg DM), dOM (up to 2.3 percentage point (pp)) and ME (up to 0.3 MJ/kg DM), but these differences were overall small. More considerable effects of pelleting were observed for uCP, which was increased in all compound feeds except the two with the highest CP concentrations. The ID<sub>RUP</sub> was lower in most compound feeds following pelleting (up to 15 pp). Pelleting also affected CP fractions in a non-systematic way. Overall, the effects of pelleting were not considerable, which could be because pelleting conditions were mild. Our third objective was to compare in situ ruminal CP degradation (ED<sub>IN\_SITU</sub>) of compound feeds with ED using two prediction methods based on CP fractions. ED<sub>IN\_SITU</sub> reference data were obtained from a companion study using the same feeds. Prediction accuracy of ED<sub>IN\_SITU</sub> and ED<sub>CNCPS</sub> was variable and depended on the compound feed and prediction method. However, future studies are needed as to date not enough data are published to draw overall conclusions for the prediction of ED<sub>IN\_SITU</sub> from CP fractions.*

**Keywords:** additivity, associative effects, *in situ* prediction, mixed feed, interaction

of organic matter (**dOM**) and metabolisable energy (**ME**), measuring gas production (**GP**) by the Hohenheim gas test (**HGT**), as described by Menke and Steingass (1988), is an established assay. An extension of this method known as extended HGT (**eHGT**; Steingass and Südekum, 2013) can be used to estimate the utilisable CP at the duodenum (**uCP**), which is the basis for the calculation of metabolisable protein used in the German protein evaluation system for cows (GfE, 2001). Calsamiglia and Stern (1995) developed a three-step method for estimating the intestinal digestibility of rumen-undegraded protein (**ID<sub>RUP</sub>**). These *in vitro* methods involve the use of ruminally fistulated animals as donors of rumen fluid. Sniffen *et al.* (1992) described a rapid CP fractionation method to be part of the Cornell Net Carbohydrate and Protein System (**CNCPS**). Therein, the CP in a feedstuff is separated into fractions by measuring N solubility. In an experiment by Chrenková *et al.* (2014), CP fractions were correlated with ruminal effective CP degradation (**ED**) values determined *in situ*. The CP fractions can be used to estimate ED values, which were found to correlate well with the ED values determined *in situ* (Shannak *et al.*, 2000). Additivity of feeding values of forages or mixes of forages and concentrates has been investigated utilising GP (Sandoval-Castro *et al.*, 2002; Robinson *et al.*, 2009; Niderkorn *et al.*, 2011) and uCP (Zhao *et al.*, 2005). However, to our knowledge, there has been no research on the additivity of **ID<sub>RUP</sub>**. Also, comprehensive data on additivity of multiple single feeds in a compound feed are not available.

Compound feeds for cattle are often in pellet form, and pelleting can increase the availability of CP and starch (**ST**) or increase indigestible bonds, depending on the intensity of the pelleting process (Svihus and Zimonja, 2011). In compound feeds, these effects can depend on the choice of single feeds, and hence, they should be examined over a wide range of various compound feeds. The objective of the present study was to characterise GP and the related values of dOM and ME as well as uCP, **ID<sub>RUP</sub>** and CP fractions of single feeds and the compound feeds produced with them, both in mash and pellet form. Three hypotheses were developed:

(**DDGS**), maize gluten, wheat bran and sugar beet pulp. Between five and seven single feeds were included in each compound feed in different concentrations. Compound feeds were produced in mash and pellet form using standard industrial processes in the feed mill of RKW-Kehl (Kehl, Germany). Production and analysed nutrient concentrations and particle size distribution of all feeds were detailed previously (Grubješić *et al.*, 2019). Targeted CP concentrations were achieved in all compound feeds. Crude protein, ash, ether extract (**EE**), NDF assayed with a heat stable amylase and expressed exclusive of residual ash (**aNDFom**) and ADF expressed exclusive of residual ash (**ADFom**) did not differ more than one percentage point (**pp**) between calculated concentrations from single feeds and analysed concentrations in mash compound feeds.

#### *Gas production kinetics, metabolisable energy and digestibility of organic matter*

*In vitro* GP kinetics were measured using HGT following the procedure described by Seifried *et al.* (2016). Approximately 200 ± 5 mg of feed ground through a 1-mm sieve was transferred into graded glass syringes (100 ml volume). Fresh rumen fluid was obtained from two rumen-fistulated Jersey cows, one lactating and one not lactating. The lactating cow was provided a ration consisting of (on DM basis) 41.3% concentrate mix, 20.0% maize silage, 16.3% meadow hay, 15.0% grass silage, 3.6% barley straw, 2.4% mineral mix and 1.4% rapeseed meal. The other cow was provided a ration consisting of 35.4% maize silage, 35.4% grass silage, 24.6% meadow hay, 3.2% barley straw, 1.0% mineral mix and 0.4% urea. Cows had *ad libitum* access to feed.

The rumen fluid obtained from the two cows was mixed to a 1 : 1 ratio, filtered through two layers of cheesecloth, and a reduced buffer solution was added. Syringes were pre-warmed to 39°C before 30 ml of buffer-rumen fluid mix was poured into each syringe under constant CO<sub>2</sub> flow. Each feed was included in five separate HGT runs with two replicated syringes per feed in each run. Additionally, each run contained three syringes without feed samples (blanks) and three syringes with a concentrate standard feed. Cumulative GP was recorded after 2, 4, 6, 8, 12, 24, 48 and 72 h of incubation at 39°C under constant rotation.

## Additivity of single feeds to compound feeds

- (a) Maize, wheat, barley, faba beans, maize gluten and wheat bran according to Krieg *et al.* (2017):

$$\text{ME (MJ/kg DM)} = 0.9065 \cdot (1.681 + 0.157\text{GP}_{24} + 0.0084\text{CP} + 0.0220\text{EE} - 0.0081\text{ash}) \quad (3)$$

- (b) Non-cereal feeds (soya beans, soya bean meal, rapeseed meal, sunflower meal, DDGS and sugar beet pulp) according to Menke and Steingass (1988):

$$\text{ME (MJ/kg DM)} = 1.06 + 0.157\text{GP}_{24} + 0.0084\text{CP} + 0.0220\text{EE} - 0.0081\text{ash} \quad (4)$$

- (c) Compound feeds according to GfE (2009):

$$\text{ME (MJ/kg DM)} = 7.17 - 0.01171\text{ash} + 0.00712\text{CP} + 0.01657\text{EE} + 0.00200\text{ST} + 0.00202\text{ADFom} + 0.06463\text{GP}_{24} \quad (5)$$

In equations (2) to (5), CP, ST, EE, ash and ADFom are expressed in g/kg DM.

### Utilisable CP at the duodenum

The eHGT method described by Steingaß and Südekum (2013) was used to estimate uCP and was conducted according to Westreicher-Kristen *et al.* (2015). Some former studies using this approach used the term 'modified HGT' and the abbreviation 'modHGT'. However, the term 'extended HGT' and the abbreviation 'eHGT' may be more appropriate as this method is not a real modification of the original HGT but an extension (measuring  $\text{NH}_3\text{-N}$  after incubation) and can be connected with  $\text{GP}_{24}$  measurement to estimate dOM and ME. Samples were incubated similarly to those in the HGT method described above. Donor cows had *ad libitum* access to a ration consisting of (on DM basis) 25.8% concentrate mix, 24.3% grass silage, 24.3% maize silage, 17.0% hay, 4.4% rapeseed meal, 2.2% barley straw and 2.0% mineral mix. Samples were incubated twice for different times (8 and 24 h), and a standard concentrate

where  $N_{\text{sample}}$  is the amount of N from the feed sample (mg),  $\text{NH}_3\text{-N}_{\text{sample}}$  and  $\text{NH}_3\text{-N}_{\text{blank}}$  are the  $\text{NH}_3\text{-N}$  content of feed samples and blank incubation residues (mg) and weight is the weight of feed sample inserted into the glass syringe (mg DM). Effective uCP was estimated for theoretical ruminal passage rates ( $k$ ) of 5 and 8%/h by plotting uCP values ( $y$ ) against the natural logarithm of the incubation time ( $x$ ) in a linear regression model and calculating the function values of  $\ln(20)$  and  $\ln(12.5)$ , respectively (Steingaß and Südekum, 2013).

### Intestinal digestibility of rumen-undegraded protein

The three-step enzymatic method of Calsamiglia and Stern (1995) was used to determine  $\text{ID}_{\text{RUP}}$ . Samples of single and compound feeds were ground through a 2-mm screen. The first step was a 16-h *in situ* incubation in the rumen, and this was conducted using three rumen-fistulated Jersey cows, following the procedure described in Seifried *et al.* (2016). A minimum of 60 mg of residual N per feed was accumulated for subsequent *in vitro* simulation of digestibility in the abomasum and duodenum. Two or three samples per feed containing 15 mg of residual N were incubated utilising 10 ml HCl (0.1 N,  $\text{pH} = 1.9$ ), pepsin (1 g/l, Sigma P-7012; Sigma, St. Louis, MO, USA) and pancreatin solution (0.5 M  $\text{KH}_2\text{PO}_4$  buffer standardised at  $\text{pH} 7.8$  containing 50 ppm of thymol and 3 g/l of pancreatin, Sigma P-7545; Sigma). Trichloroacetic acid was added to stop enzymatic action and precipitate undigested proteins. Samples were centrifuged at 15 000 g for 25 min. Supernatants were analysed for soluble N by the Kjeldahl method (VDLUFA, 2007). Finally,  $\text{ID}_{\text{RUP}}$  was calculated as follows:

$$\text{ID}_{\text{RUP}} (\%) = (N_{\text{soluble}} / N_{\text{incubated}}) \cdot 100 \quad (7)$$

where  $N_{\text{soluble}}$  is the amount of soluble N determined *in vitro* (mg) and  $N_{\text{incubated}}$  is the total N that was incubated with pepsin and pancreatin (mg).

### CP fractionation

Crude protein fractions were estimated according to the CNCPS (Sniffen *et al.*, 1992): fraction A represented the non-protein N, fraction B the true protein and containing

where  $ED_{CNCPS}$  is ED calculated from CP fractions, A and B1 to B3 are determined CP fractions, Prot-B1 to Prot-B3 are table values for ruminal degradation rates of CP fractions and  $k$  is the ruminal passage rate (5 or 8%/h). The ruminal degradation rate of faba beans was not reported in Fox *et al.* (2003), and thus, the value for lupins was used instead as they both belong to the legume family and show similar ED values (Goelema *et al.*, 1998). Table values of ruminal degradation rates of compound feeds were not available; therefore, they were calculated from values of the respective single feeds. This calculation included weighting contributions of CP of single feeds to the total CP of the respective compound feed. Ruminal degradation rates were then used together with determined CP fractions to estimate the observed  $ED_{CNCPS}$  for mash and pelleted compound feeds.

An alternative prediction equation (Shannak *et al.*, 2000) based on CP fractions and NDF was used to estimate RUP of compound feed as follows:

$$\begin{aligned} RUP_5 \text{ or } RUP_8 (\text{g/kg CP}) = & \beta_0 + \beta_1 \text{ CPPNDF} \\ & + \beta_2 (\text{CP} \times \text{B2}) + \beta_3 (\text{CP} \times \text{C}) + \beta_4 (\text{CP} (\text{A} + \text{B1})) \\ & + \beta_5 (\text{CP} \times \text{C}^2) + \beta_6 (\text{PNDF} \times \text{B1}) \\ & + \beta_7 ((\text{B3} + \text{C}) \text{B2}) + e \end{aligned} \quad (9)$$

where  $RUP_5$  or  $RUP_8$  are RUP values for rumen outflow rates of  $k=5$  and 8%/h, respectively. The CPPNDF is the CP concentration in PNDF (NDF determined by manual filtration on paper) and all nutrients are given as g/kg DM, whereas A, B and C fractions are given as g/kg CP. Instead of CPPNDF and PNDF, the CP concentration in aNDF as well as aNDF was determined in the present study by the conventional method (VDLUF, 2007). The general form of the equation is identical for  $RUP_5$  and  $RUP_8$ , and the parameter estimated of  $\beta_0$  to  $\beta_7$  is given in Shannak *et al.* (2000). The  $ED_{CNCPS}$  was then calculated from RUP values for the given rumen outflow as:

$$ED_{CNCPS} (\% \text{ of CP}) = (1000 - \text{RUP})/10 \quad (10)$$

values of single feeds were determined according to either equations (3) or (4) depending on the feed group or alternatively equation (5) for all single feeds.

#### Statistical analyses

Calculated and observed values of mash compound feeds, and values of mash and pellet compound feeds, were regressed using procedure REG (version 9.4 of SAS system for Windows SAS Institute, Cary, NC, USA). The REG procedure was also used to calculate if slopes and intercepts were significantly different from 1 and 0, respectively, by determination of 95% CI to detect possible associative effects.

## Results

### *In vitro ruminal fermentation and feeding values of single feeds*

Overall, ruminal fermentation characteristics and feeding values of single feeds varied widely (Table 1). The highest  $GP_{24}$  was found in maize (81 ml/200 mg DM) and the lowest in sunflower meal (36 ml/200 mg DM). The highest dOM was found in maize and wheat (96%) and the lowest in sunflower meal (65%). The highest ME was found in soya beans (16.0 MJ/kg DM) followed by maize (14.5 MJ/kg DM) and wheat (14.2 MJ/kg DM) and the lowest in sunflower meal (9.4 MJ/kg DM). The uCP concentration varied between 144 g/kg DM in wheat bran and 279 g/kg DM in soya bean meal for  $k=5\%/h$ , and between 158 g/kg DM in maize and wheat bran and 356 g/kg DM in soya bean meal for  $k=8\%/h$ . The  $ID_{RUP}$  ranged between 18% in wheat bran and 83% in maize and soya bean meal. All CP fractions varied widely between single feeds. This was reflected in the high variability of  $ED_{CNCPS}$  values.

### *Additivity of fermentation characteristics and feeding values*

Calculated and observed ruminal fermentation characteristics and nutritional values of the mash compound feeds are presented in Table 2. Estimation of  $bGP$ ,  $GP_{24}$  and dOM was precise as indicated by the slope of the regression lines (close to 1) and the high  $R^2$  values (Table 3). Observed cGP differed numerically (0.3 to 0.7 pp) from calculated values,

Con	GP <sub>24</sub> /200 mg DM)	DOM	ME (MJ/kg DM)	uCP		ID <sub>RUP</sub> (%)	CP fractions					ED <sub>CNCPS</sub>	
				k = 5%/h (g/kg DM)	k = 8%/h (g/kg DM)		A (%)	B1 (%)	B2 (%)	B3 (%)	C (%)	k = 5%/h (%)	k = 8%/h (%)
81		96	14.5	163	158	83	12.5	5.9	66.6	14.9	0.0	65	55
78		96	14.2	180	198	35	9.8	22.2	63.5	4.4	0.0	80	73
75		92	13.6	162	175	47	11.4	16.4	62.3	5.0	5.0	76	69
46		79	16.0	227	282	71	4.5	4.8	82.7	6.4	1.6	56	45
51		92	13.3	279	356	83	2.9	8.4	83.8	3.6	1.2	72	62
45		78	11.3	241	280	26	7.1	13.4	62.5	10.2	6.8	73	65
36		65	9.4	166	205	46	8.3	27.3	54.6	5.9	3.9	76	69
63		90	12.9	202	224	32	12.7	42.3	38.7	4.2	2.1	80	75
44		73	11.9	275	296	75	20.6	0.0	44.1	11.8	23.6	55	50
54		78	11.0	188	196	59	48.5	6.2	33.8	8.6	2.9	79	75
52		72	10.9	144	158	18	12.5	21.1	49.4	13.6	3.4	80	74
74		90	13.2	182	199	67	39.9	0.0	29.2	24.8	6.2	82	78

Con, GP<sub>24</sub> = corrected gas production at 24 h; DOM = digestibility of organic matter; ME = metabolisable energy; uCP = utilisable CP for ruminal passage rates (k) of 5 and 8%/h; ID<sub>RUP</sub> = ruminal passage rates; CP fractions = crude protein fractions: A = non-protein nitrogen; B1 = rapidly degradable true protein; B2 = moderately degradable true protein; B3 = slowly degradable true protein, determined using Cornell Net Carbohydrate and Protein System (CNCPS); ED<sub>CNCPS</sub> = effective protein degradation for ruminal passage rates of 5 and 8%/h.

## Additivity of single feeds to compound feeds

Observed ID<sub>RUP</sub> in compound feeds differed from calculated values between 0 and 11 pp. Regression analysis between calculated and observed ID<sub>RUP</sub> values showed an  $R^2$  value of 0.64 and relatively large CIs for the slope and intercept, respectively.

Conformity between calculated and observed CP fractions depended on the specific fraction and the compound feed type. Confidence interval of the slope did not include the value of 1 only for B3 (CI = 1.22 to 2.18), even though the  $R^2$  value for this parameter was high ( $R^2 = 0.93$ ).

Results of the regression analysis of calculated and observed ED<sub>CNCPS</sub> values showed small accuracy ( $R^2$  of 0.36 and 0.43 for  $k = 5$  and  $8\%/h$ , respectively). However, the slope values included 1 and intercept values included 0 and numerical differences were in most cases not even detectable (Table 2).

### Effects of pelleting on ruminal fermentation characteristics and feeding value of compound feeds

Differences between mash and pellet compound feeds in GP<sub>24</sub> did not exceed 3 ml/200 mg DM, 3 pp in dOM and 0.3 MJ ME/kg DM (Table 2). However, based on CI ranges (Table 3), the results indicated that pelleting did affect GP characteristics. The slopes and the intercepts for bGP, cGP, GP<sub>24</sub>, dOM and ME were all significantly different from 1 to 0, respectively, even though the  $R^2$  value was 0.93 or higher. Pelleting numerically increased uCP in compound feeds with lower CP concentration and decreased uCP in compound feeds with higher CP concentration. The slopes and intercepts were significantly different from 1 and 0, respectively, with considerable differences in  $R^2$  values between  $k = 5\%/h$  ( $R^2 = 0.38$ ) and  $k = 8\%/h$  ( $R^2 = 0.83$ ). Pelleting decreased estimated ID<sub>RUP</sub> in most compound feeds, with a maximum of 15 pp in compound feed 3. Pelleting increased estimated ID<sub>RUP</sub> only in compound feed 1, but the difference was negligible (2 pp). Although the CI for the slope of ID<sub>RUP</sub> included 1 and the  $R^2$  value was high ( $R^2 = 0.92$ ), the intercept was significantly different from 0. Pelleting did not systematically affect CP fractions in compound feeds (Table 3). Pelleting reduced the ED<sub>CNCPS</sub> in most compound feeds slightly (up to 3 pp) for both  $k = 5$  and  $8\%/h$ .



**Table 2.** *In vitro* fermentation characteristics and feeding value of mash and pellet compound feeds and values calculated from single feeds

Compound feed	<i>In vitro</i> gas production			DOM	uCP		ID <sub>RUP</sub>	CP fractions					ED <sub>CNCPS</sub>	
	bGP (ml/200 mg DM)	cGP (%/h)	GP <sub>24</sub> (ml/200 mg DM)		k = 5%/h (g/kg DM)	k = 8%/h (g/kg DM)		A (%)	B1 (%)	B2 (%)	B3 (%)	C (%)	k = 5%/h (%)	k = 8%/h (%)
1 Calculated	77	7.0	73	92	181	193	63	12.9	12	66.3	7.9	0.9	75	67
Mash	81	7.3	74	93	173	188	71	17.1	7.1	68.0	7.8	0.0	75	67
Pellet	77	8.4	71	91	192	212	73	19.4	3.0	69.9	7.7	0.0	74	66
2 Calculated	69	7.9	66	87	190	205	46	16.5	11.8	54.5	10.8	6.4	76	69
Mash	71	8.6	65	86	177	195	55	23.2	3.9	55.3	14.1	3.5	76	69
Pellet	70	9.2	65	86	185	204	48	21.7	5.9	54.9	14.0	3.5	76	69
3 Calculated	68	8.2	62	84	175	193	44	17.3	21.3	51.3	6.8	3.2	77	70
Mash	67	8.3	61	83	173	188	49	18.1	19.5	52.9	6.3	3.2	77	70
Pellet	67	9.1	62	83	178	198	34	17.4	16.9	56.3	6.3	3.2	76	69
4 Calculated	72	8.2	66	89	192	216	59	9.3	15.3	65.3	6.0	4.1	71	62
Mash	73	7.5	67	90	186	210	70	8.8	14.7	67.9	5.7	2.9	71	62
Pellet	72	8.5	66	89	190	214	63	10.1	15.6	65.5	5.8	2.9	72	63
5 Calculated	68	7.6	63	87	200	222	49	9.3	18.6	59.8	7.7	4.6	73	66
Mash	69	8.1	62	86	191	209	58	10.1	18.9	57.8	7.9	5.3	73	66
Pellet	68	9.0	63	86	190	213	51	9.0	17.8	62.2	5.5	5.5	72	65
6 Calculated	63	8.7	58	83	196	225	49	9.3	19.0	61.1	7.3	3.3	73	66
Mash	64	8.4	59	84	189	213	49	9.0	21.1	60.0	7.4	2.5	74	66
Pellet	66	8.9	61	85	194	217	44	9.5	19.9	59.9	5.4	5.4	71	64
7 Calculated	55	8.9	54	81	199	233	50	9.3	14.1	66.4	7.1	3.0	73	65
Mash	59	9.2	54	81	195	226	53	9.4	14.4	64.9	9.0	2.2	73	65
Pellet	59	9.5	55	81	188	220	41	10.5	12.6	67.5	7.1	2.4	72	64
8 Calculated	62	8.7	57	85	203	240	60	7.1	17.9	67.6	5.0	2.4	73	65
Mash	64	8.4	59	87	199	237	56	9.4	18.4	66.0	4.2	2.1	73	65
Pellet	63	9.2	59	86	198	233	48	7.7	17.8	68.0	4.3	2.1	72	64


bGP = potential gas production; cGP = rate of gas production; GP<sub>24</sub> = corrected gas production at 24 h; DOM = digestibility of organic matter; uCP = utilisable CP for ruminal passage rates (*k*) of 5 and 8%/h; ID<sub>RUP</sub> = intestinal digestibility of rumen-undegraded protein; CP fractions = crude protein fractions: A = non-protein nitrogen; B1 = rapidly degradable true protein; B2 = moderately degradable true protein; B3 = slowly degradable true protein, C = undegradable and indigestible true protein, determined using Cornell Net Carbohydrate and Protein System (CNCPS); ED<sub>CNCPS</sub> = Effective protein degradation for ruminal passage rates of 5 and 8%/h, calculated using Fox *et al.* (2003). Metabolisable energy (ME) values are presented in Figure 1.

**Table 3.** Results of simple linear regressions for *in vitro* fermentation characteristics and feeding values of compound feeds


	Calculated v. observed						Mash v. pelleted					
	Slope	Slope CI	Intercept	Intercept CI	R <sup>2</sup>	RMSE	Slope	Slope CI	Intercept	Intercept CI	R <sup>2</sup>	RMSE
<i>In vitro</i> gas production												
<i>b</i> GP	0.98	0.74 to 1.21	3.58	−11.95 to 19.11	0.95	1.70	0.79	0.64 to 0.93	13.80	3.87 to 23.72	0.97	1.06
<i>c</i> GP	0.68	0.03 to 1.33	2.67	−2.65 to 7.98	0.52	0.44	0.59	0.43 to 0.75	4.14	2.82 to 5.47	0.93	0.10
GP <sub>24</sub>	0.96	0.78 to 1.14	2.72	−8.52 to 13.95	0.97	1.19	0.82	0.67 to 0.98	11.27	1.63 to 20.91	0.97	1.00
dOM	1.10	0.73 to 1.40	−4.95	−33.43 to 23.53	0.91	1.25	0.77	0.59 to 0.94	19.84	4.66 to 35.02	0.95	0.74
ME	− <sup>1</sup>	—	—	—	—	—	0.71	0.54 to 0.87	3.82	1.61 to 6.03	0.95	0.07
uCP												
<i>k</i> = 5%/h	0.97	0.61 to 1.33	−1.05	−70.00 to 67.89	0.88	3.73	0.37	−0.11 to 0.85	120.09	31.12 to 209.07	0.38	5.16
<i>k</i> = 8%/h	0.97	0.77 to 1.17	−1.26	−45.18 to 42.66	0.96	3.89	0.53	0.29 to 0.77	103.00	53.24 to 152.76	0.83	4.53
ID <sub>RUP</sub>	0.95	0.23 to 1.66	7.86	−30.08 to 45.80	0.64	5.53	1.42	0.99 to 1.85	−31.27	−56.26 to −6.28	0.92	3.94
CP fractions												
A	1.37	0.79 to 1.94	−24.19	−92.26 to 43.88	0.85	23.09	0.95	0.69 to 1.20	7.15	−28.95 to 43.25	0.93	15.30
B1	1.65	0.94 to 2.35	−119.87	−237.17 to −2.56	0.84	26.55	0.94	0.64 to 1.24	−2.20	−49.96 to 45.55	0.91	20.23
B2	0.94	0.64 to 1.23	38.34	−144.37 to 221.05	0.91	19.22	0.88	0.52 to 1.24	88.96	−131.32 to 309.24	0.86	22.79
B3	1.70	1.22 to 2.18	−46.79	−82.89 to −10.69	0.93	8.71	0.94	0.57 to 1.32	−3.55	−34.68 to 27.58	0.86	12.03
C	0.71	0.13 to 1.29	2.36	−19.67 to 24.40	0.60	10.21	0.98	0.31 to 1.65	4.60	−15.84 to 25.04	0.68	10.83
ED <sub>CNCPS</sub>												
<i>k</i> = 5%/h	1.24	−0.41 to 2.89	−12.35	−134.36 to 109.66	0.36	3.40	0.63	0.14 to 1.11	31.83	−7.02 to 70.68	0.62	2.08
<i>k</i> = 8%/h	1.41	−0.20 to 3.02	−20.20	−126.68 to 86.28	0.43	4.22	0.65	0.19 to 1.10	28.54	−4.71 to 61.79	0.67	2.55

*b*GP = potential gas production; *c*GP = rate of gas production; GP<sub>24</sub> = corrected gas production at 24 h; dOM = digestibility of organic matter; ME = metabolisable energy; uCP = utilisable CP for ruminal passage rates (*k*) of 5 and 8%/h; ID<sub>RUP</sub> = intestinal digestibility of rumen-undegraded protein; CP fractions = crude protein fractions: A = non-protein nitrogen; B1 = rapidly degradable true protein; B2 = moderately degradable true protein; B3 = slowly degradable true protein, C = undegradable and indigestible true protein, determined using Cornell Net Carbohydrate and Protein System (CNCPS); ED<sub>CNCPS</sub> = effective protein degradation for ruminal passage rates of 5 and 8%/h, calculated using Fox *et al.* (2003). Observed values refer to values for compound feeds in mash form.

<sup>1</sup>Simple linear regressions of calculated and observed ME values are presented in Figure 1.



**Figure 1.** Comparison of calculated and observed metabolisable energy (ME) values of compound feeds using an *in vitro* ruminal fermentation technique. The ME values of compound feeds were calculated from ME values of single feeds that were determined according to the equations of: (a) Krieg *et al.* (2017) and Menke and Steingass (1988), respective of the feed group; or (b) GfE (2009) for all single feeds. The dotted line represents the angle bisector.





concentrations are often specific to single feeds or groups of feeds because this is associated with high prediction accuracy. However, it does not necessarily hold true across different feed groups and also for feed mixtures (Menke and Steingass, 1988). This caused issues when values for single feeds and related compound feeds were compared in the present study. We argue that the conclusion that additivity does not exist is misleading because it is an artefact of using different equations for different groups of feeds. When equation (5) was used to calculate the ME for all feeds (single and compound), the differences between calculated and observed ME became negligible (Figure 1).

Gas production techniques are underutilised for estimation of potential interactions between feeds (D'Mello, 2000) and energy evaluation, and previous research on additivity has most commonly focused on concentrate–forage mixes. Robinson *et al.* (2009) noted associative effects (15% to 25%) in mixture of alfalfa hay, barley grain, maize silage and soya bean meal using HGT in early phase of incubation, while they disappeared later. Similarly, Arhab *et al.* (2010) used mixtures of triticale and barley with a commercial concentrate supplement and found significant differences only in GP up to 8 h. In the present study, the discrepancy between calculated and observed GP at multiple time points can be considered to be negligible with a maximal difference of 2 ml/200 mg DM after 2 and 4 h of incubation. The observed GP<sub>24</sub> was similar to the calculated ones and was considered to be additive. Since dOM and ME values are derived from GP<sub>24</sub> (Menke and Steingass, 1988), this is an important finding and we presume that the differences between calculated and observed ME of compound feeds were caused only by the choice of ME equation. For the estimation of dOM values, the SD of regression residuals ( $s_{y,x}$ ) is 3.07% (Menke and Steingass, 1988). The  $s_{y,x}$  for estimation of ME values depends on the regression equation used, with a  $s_{y,x}$  of 2.92 MJ/kg DM (Menke and Steingass, 1988) or a RMSE value of 1.98 MJ/kg DM (GfE, 2009) for equations (4) and (5), respectively. The RMSE between calculated and observed dOM and ME (equation (5)) of the compound feeds of the present study was 1.15% and 0.09 MJ/kg DM, respectively, and therefore markedly lower compared to the  $s_{y,x}$  and the

should be interpreted with caution and reference to their biological and practical relevance, which seems negligible. This contrasts with the findings of Zhao *et al.* (2005), who reported higher differences between calculated and observed uCP values in an experiment using 16 single feeds and 19 mixtures. They noted statistically significant and non-systematic differences between calculated and observed uCP values. However, the authors mentioned the possibility of incomplete incubation of some feeds due to incomplete mixing with the incubation liquid (Zhao *et al.*, 2005). Such an effect was avoided in the present study by the constant motion of the rotary incubator.

Calsamiglia and Stern (1995) highlighted the importance of ID<sub>RUP</sub> for evaluating feed protein. GfE (2001) and NRC (2001) assumed a constant ID<sub>RUP</sub> value of 80% for all feeds. In the CNCPS, ID<sub>RUP</sub> was assumed to be 100% for CP fractions A, B1 and B2, 80% for fraction B3 and zero for fraction C (Fox *et al.*, 2003). A wide range of ID<sub>RUP</sub> of single and compound feeds was found in the present study. Observed ID<sub>RUP</sub> in most compound feeds was higher than calculated (up to 11 pp). Accurate estimation of ID<sub>RUP</sub> from single feeds was thus not possible for all compound feeds of the present study using the three-step method. This is underlined by the analytical tolerance of the determination of ID<sub>RUP</sub> which was set to maximal 10% relative deviation from the mean value otherwise the procedure was repeated. Relative deviation of replicates varied between 0.04 and 8.11% around the mean value for all feed samples of the present study. However, for five out of eight compound feeds, relative deviations of calculated ID<sub>RUP</sub> from observed ID<sub>RUP</sub> exceeded the value of 10%, which represents the analytical tolerance. This indicates that associative effects occurred when analysing ID<sub>RUP</sub> of compound feeds. Those interactions between single feeds in mixture can occur in any of the three steps. Calculation of the 16 h *in situ* CP degradation of compound feeds from single feeds showed better additivity than ID<sub>RUP</sub> with a slight tendency to overestimate CP degradation (1 to 7 pp). Compound feeds that showed higher deviations between calculated and observed *in situ* degradation also tended to show higher differences between the calculated and observed ID<sub>RUP</sub>. Hence, associative effects seem more pronounced during

values were higher than analytical tolerances for CP analysis (VDLUFA, 2019). This was the case for the A and B1 fraction of compound feeds 1 and 2, and the B3 fraction of compound feed 2. However, for all other CP fractions and compound feeds, the difference between the calculated and observed values of CP fractions is similar or even lower than the analytical tolerance of CP analysis. In addition, small variability between compound feeds (particularly for CP fraction C) probably lowered the accuracy of regression analysis. Consequently, interpretation of additivity for CP fractions is difficult from the results of the present study and different depending on the specific fraction and feed type. Additivity of  $ED_{CNCPS}$  was given for all compound feeds. However, the accuracy of regression analysis may be limited owing to the relatively small sample size ( $n = 8$  compound feeds) of the present study. Therefore, we recommend to examine the additivity of CP fractions of single feeds in compound feeds in further experiments.

#### *Effects of pelleting on ruminal fermentation characteristics and feeding value of compound feeds*

The second hypothesis of the present study was that the pelleting process would significantly affect GP, dOM, ME, uCP,  $ID_{RUP}$  and CP fractions of compound feeds. Based on the present results, this hypothesis can be rejected. Even though the results of statistical analysis indicated an effect of pelleting on GP and related values of ME and dOM, uCP and  $ID_{RUP}$ , the overall numerical differences were negligible.

When heat is excessively applied during the processing of compound feeds, the intestinal digestibility of protein can be reduced owing to the formation of Maillard products which can neither be fermented nor digested (Sniffen *et al.*, 1992). Any optimum of processing conditions would aim to reduce CP degradability in the rumen without affecting  $ID_{RUP}$ . The data obtained *in situ* with the same feeds as used in the present study (Grubješić *et al.*, 2019) indicated that pelleting increased rumen degradation of some compound feeds, thus resulting in less RUP entering the small intestine. However, pelleting increased the share of smaller feed particles compared with the mash feeds, which might have increased the number of feed particles leaving the bags without microbial degradation, and thus overestimated

result in chemical or physical changes of the substrate. In the present study, except for compound feed 1,  $ID_{RUP}$  decreased from 6 to 15 pp in all compound feeds by pelleting.

*In situ* incubations over 16 h were used to generate RUP for *in vitro* determination of  $ID_{RUP}$ , and results showed that degradation after 16 h increased between 1.4 and 6.4 pp in pelleted compound feeds compared to their corresponding mash feeds. It can therefore be assumed that RUP of mash feeds after *in situ* incubation contained more potentially digestible CP for the *in vitro* enzymatic steps to determine  $ID_{RUP}$ . This is underlined by the calculation of total tract digestibility (TTD) from the summation of 16 h *in situ* RUP and *in vitro*  $ID_{RUP}$  which showed that differences in TTD between mash and pelleted compound feeds ranged only between 0.2 and 2.2 pp and can therefore be considered to be negligible. The higher rumen-degraded protein of pelleted compound feeds might be attributed to a smaller particle size compared to mash feeds, as explained in the previous sections.

Pelleting did not have a large effect on CP fractions and  $ED_{CNCPS}$  values of compound feeds. Heat treatment during the pelleting process can denaturise protein fraction B2 making it insoluble, resulting in increased B2 and C fractions (Licitra *et al.*, 1996). Such an effect was not found in the present study, probably due to the temperature during pelleting not being very high.

#### *Prediction of in situ ruminal CP degradation from CP fractions*

The third hypothesis of the present study was that  $ED_{IN\_SITU}$  could be predicted using CP fractions. Based on the present results, this hypothesis is rejected. Compared with the corresponding  $ED_{IN\_SITU}$  data (Grubješić *et al.*, 2019), neither the calculation of  $ED_{CNCPS}$  using individual CP fractions and tabular values for their specific degradation rates (Fox *et al.*, 2003) nor  $ED_{CNCPS}$  using proximate nutrients and CP fractions based on regression analysis (Shannak *et al.*, 2000) showed adequate prediction accuracy for all compound feeds. However, for two (calculated according to Fox *et al.* (2003)) and three (calculated according to Shannak *et al.* (2000)) out of eight compound feeds, ED prediction with both methods was similar

## Additivity of single feeds to compound feeds

derived their prediction equations from selected proximate nutrients and CP fractions for *in situ* RUP values including 11 dairy compound feeds. Therefore, prediction of ED of compound feeds may be possible with good accuracy. Shannak *et al.* (2000) found differences between *in situ* RUP values and respective estimates of up to 79 g/kg CP; however, 8 out of 11 RUP values had differences  $\leq 50$  g/kg CP. For samples of the present study, ED<sub>IN\_SITU</sub> and ED<sub>CNCPS</sub> calculated according to Shannak *et al.* (2000) differed by up to 16 pp and hence 5 out of 8 compound feeds had differences between estimated and *in situ* RUP  $\geq 100$  g/kg CP for  $k=5$  and 8%/h. Poor estimation may result from differences in the assay details because NDF was determined by manual filtration in the study of Shannak *et al.* (2000), and authors stated that results may deviate from those obtained with the conventional NDF method which was used in the present study. Moreover, NDF values ranged between 212 and 554 g/kg DM in the 11 compound feeds of Shannak *et al.* (2000) and only between 142 and 255 g/kg DM in the present study. Shannak *et al.* (2000) also included forages and special by-products in the development of the regression equations, which is another difference to the present study. It is therefore recommended to extend the existing database. More accurate equations may be developed when covering a wider range of feedstuff groups.

## Conclusion

We conclude that, when formulating compound feeds for cattle, single feed data for GP<sub>24</sub>, dOM, ME and uCP are additive, while those for ID<sub>RUP</sub> are not. Additivity of CP fractions is dependent on the fraction and compound feed type, whereas ED<sub>CNCPS</sub> is precisely additive. The pelleting process had little effect on ruminal fermentation characteristics and feeding values of compound feeds, probably because heat exposure was moderate. Using CP fractions in the present study did not reliably predict *in situ* ruminal CP degradation of compound feeds: more studies are needed to extend the database for the development of prediction equations.

## Acknowledgements

## Ethics statement

The use of rumen-cannulated cows for rumen fluid collection was approved by the Regierungspräsidium Stuttgart, Germany, approval number A401-14 TE.

## Software and data repository resources

No data were deposited in an official repository.

## References

- Arhab R, Laadjimi K, Driss D, Djabri B and Boussebouda H 2010. Evaluation of feed mixture interactions by using *in vitro* gas production method. *Livestock Research for Rural Development* 22, Article #217.
- Calsamiglia S and Stern MD 1995. A three-step *in vitro* procedure for estimating intestinal digestion of protein in ruminants. *Journal of Animal Science* 73, 1459–1465.
- Chrenková M, Čerešňáková Z, Weisbjerg MR, Formelová Z, Poláčiková M and Vondráková M 2014. Characterization of proteins in feeds according to the CNCPS and comparison to *in situ* parameters. *Czech Journal of Animal Science* 59, 288–295.
- D'Mello JPF 2000. *Farm animal metabolism and nutrition*. CABI Pub., Scottish Agricultural College, Edinburgh, UK.
- Fox DG, Tylutki TP, Tedeschi LO, Van Amburgh ME, Chase LE, Pell AN, Overton TR and Russell JB 2003. The net carbohydrate and protein system for evaluating herd nutrition and nutrient excretion: CNCPS version 5.0, model documentation. In *Animal Science Mimeo* 213, Animal Science Department, Cornell University, Ithaca, NY, USA.
- Gesellschaft für Ernährungsphysiologie (GfE) 2001. Energie-und Nährstoffbedarf landwirtschaftlicher Nutztiere. Nr. 8 Empfehlungen zur Energie-und Nährstoffversorgung der Milchkühe und Aufzuchtrinder. DLG-Verlag Frankfurt am Main, Germany.
- Gesellschaft für Ernährungsphysiologie (GfE) 2009. New equations for predicting metabolisable energy of compound feeds for cattle. In *Proceedings of the Society of Nutrition Physiology*, 10–12 March 2009, Göttingen, Germany, pp. 143–146.
- Gesellschaft für Ernährungsphysiologie (GfE) 2017. Opinion on the indispensability of animal experiments in animal nutrition research and suitability of alternative methods. Retrieved on 26 January 2020, from <https://gfe-frankfurt.de/2017/02/12/tierversuche/>
- Goelema JO, Spreeuwenberg MAM, Hof G, van der Poel AFB and Tamminga S 1998. Effect of pressure toasting on the rumen degradability and intestinal digestibility of whole and broken peas, lupins and faba beans and a mixture of these feedstuffs. *Animal Feed Science and Technology* 76, 35–50.
- Grubješić G, Titze N, Krieg J and Rodehutscord M 2019. Determination of *in situ* ruminal crude protein and starch degradation values of compound feeds from single feeds. *Archives of Animal Nutrition* 73, 414–429.
- Krieg J, Seifried N, Steingass H and Rodehutscord M 2017. *In situ* and *in vitro* ruminal starch degradation of grains from different rye, triticale and barley genotypes. *Animal* 11, 1745–1753.

## Grubješić, Titze, Krieg and Rodehutschord

Seifried N, Steingass H, Hoffmann N and Rodehutschord M 2016. *In situ* starch and crude protein degradation in the rumen and *in vitro* gas production kinetics of wheat genotypes. *Journal of Animal Physiology and Animal Nutrition* 101, 779–790.

Shannak S, Südekum KH and Susenbeth A 2000. Estimating ruminal crude protein degradation with *in situ* and chemical fractionation procedures. *Animal Feed Science and Technology* 85, 195–214.

Sniffen CJ, O'Connor JD, Van Soest PJ, Fox DG and Russell JB 1992. A net carbohydrate and protein system for evaluating cattle diets: II. Carbohydrate and protein availability. *Journal of Animal Science* 70, 3562–3577.

Steingass H and Südekum KH 2013. Proteinbewertung beim Wiederkäuer—Grundlagen, analytische Entwicklungen und Perspektiven. *Übersichten zur Tierernährung* 41, 51–73.

Svihus B and Zimonja O 2011. Chemical alterations with nutritional consequences due to pelleting animal feeds: a review. *Animal Production Science* 51, 590–596.

Titze N, Krieg J, Steingass H and Rodehutschord M 2018. Variation of lupin protein degradation in ruminants studied *in situ* and using chemical protein fractions. *Animal* 13, 709–717.

Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten (VDLUFA) 2007. *Handbuch der Landwirtschaftlichen Versuchs- und Untersuchungsmethodik Bd. III: Die chemische Untersuchung von Futtermitteln*. VDLUFA-Verlag, Darmstadt, Germany.

Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten (VDLUFA) 2019. *Analysenspielräume (ASR). Version 12* Retrieved on 15 January 2020, from [https://www.vdlufa.de/Dokumente/Fachgruppen/FG6/ASR\\_Version\\_12\\_2019.pdf](https://www.vdlufa.de/Dokumente/Fachgruppen/FG6/ASR_Version_12_2019.pdf)

Westreicher-Kristen E, Steingass H and Rodehutschord M 2015. Estimation of utilisable crude protein at the duodenum of dried distillers' grains with solubles using a modified gas test. *Archives of Animal Nutrition* 69, 351–365.

Zhao GY, Li YX, Ren JB, Li YJ and Guo DS 2005. The influence of associative effects on the *in vitro*-estimated utilizable crude protein (uCP) of feeds for ruminants. *Archives of Animal Nutrition* 59, 149–154.